

AD-A065770

TECHNICAL
LIBRARY

IMPROPER FERROELECTRICS

Structural Studies of LiNH_4SO_4

Paul M. Raccach
James S. Kouvel
Rastko C. Maglic

Department of Physics
University of Illinois at Chicago Circle
Chicago, Illinois 60680

December 1978

FINAL TECHNICAL REPORT

1 June 1977 - 31 May 1978

prepared for

Night Vision & Electro-Optics Laboratories
USAECOM
Fort Belvoir, Virginia 22060

ARPA Contract No. DAAK70-77-C-0149

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) IMPROPER FERROELECTRICS Structural Studies of LiNH_4SO_4		5. TYPE OF REPORT & PERIOD COVERED Final Technical Report 1 June 1977 - 31 May 1978
		6. PERFORMING ORG. REPORT NUMBER.
7. AUTHOR(s) Paul M. Raccah James S. Kouvel Rastko C. Maglic		8. CONTRACT OR GRANT NUMBER(s) ARPA DAAK 70-77-C-0149
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics University of Illinois at Chicago Circle Chicago, Illinois 60680		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Night Vision & Electro-Optics Labs., USAECOM Fort Belvoir, Virginia 22060		12. REPORT DATE December 1978
		13. NUMBER OF PAGES 17
14. MONITORING AGENCY NAME & ADDRESS (If different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Distribution list appended to Report.		
17. DISTRIBUTION STATEMENT (of this abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Improper Ferroelectrics; Lithium Ammonium Sulphate; Neutron Diffraction; X-Ray Diffraction		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The room temperature crystal structure of LiNH_4SO_4 , complete with the positions of the hydrogen ions, has been determined from neutron diffraction measurements. X-ray diffraction study has shown that the abrupt ferroelectric transition of LiNH_4SO_4 near 11°C involves a doubling of the orthorhombic unit cell normal to the polar axis. The phonon dispersion in this material above the transition temperature was explored by means of inelastic neutron scattering, but no soft phonon mode associated with the structural transition could be found.		

SUMMARY

The room temperature crystal structure of LiNH_4SO_4 , complete with the positions of the hydrogen ions, has been determined from neutron diffraction measurements. X-ray diffraction study has shown that the abrupt ferroelectric transition of LiNH_4SO_4 near 11°C involves a doubling of the orthorhombic unit cell normal to the polar axis. The phonon dispersion in this material above the transition temperature was explored by means of inelastic neutron scattering, but no soft phonon mode associated with the structural transition could be found.

I. INTRODUCTION

The class of ferroelectrics known as improper, whose long-range order parameter is a crystal distortion rather than the electric polarization, is currently of considerable scientific and technological interest. According to present understanding, the crystal distortion of such a ferroelectric results from the freezing-in of a non-polar (Brillouin-zone edge) phonon mode, produces a spontaneous ferroelectric polarization.[1,2] The dielectric constant of an improper ferroelectric, unlike that of a proper ferroelectric, does not diverge near the transition temperature, yet its pyroelectric coefficient can remain substantially high. This combination of properties can express itself in a large figure-of-merit for vidicon thermal-imager applications.[3]

The close interplay between the structural and electrical behavior of an improper ferroelectric makes it particularly important that the measured properties of any such material of interest be complemented by detailed investigations of its crystal structure and its static and dynamic structural changes with temperature.

The present studies have focused on the structure of LiNH_4SO_4 , an improper ferroelectric compound whose thermodynamic and optical properties have recently been explored quite extensively.[3-8] This compound undergoes a ferroelectric order-disorder transition at $T_{c1} \approx 186^\circ\text{C}$, above which, in the paraelectric phase, the temperature dependence of its dielectric constant shows

a marked departure from the Curie-Weiss law, indicative of "improper" behavior. At $T_{c2} \approx 11^\circ\text{C}$, it transforms abruptly into a different low-temperature ferroelectric state. Thus, between T_{c1} and T_{c2} , the material is in an intermediate ferroelectric phase, which is of particular application interest near room temperature.[3]

The room-temperature crystal structure of LiNH_4SO_4 has been previously found to belong to the orthorhombic space group $P2_1\text{cn}$. [9] However, since the crystal structure determination was made by x-ray diffraction, the positions of the hydrogen ions could not be located. We have corrected this situation by means of a neutron diffraction study of a LiNH_4SO_4 crystal. From our determination of its crystal structure, complete with hydrogen-ion positions, it is now possible to calculate the spontaneous electric polarization of the room-temperature phase of this material on the basis of various model assumptions. Some preliminary calculations are presented, together with our diffraction results for the structure, in Section II of this report.

The structural nature of the transition in LiNH_4SO_4 , when it is cooled through its lower critical temperature T_{c2} , has recently been explored in an optical Raman study [8], from which it was claimed that the transition involved a doubling of the primitive unit cell. The experimental evidence for this claim, however, cannot be said to be definitive. We have therefore investigated this question with x-ray diffraction measurements, and our results

described in Section III show unambiguously that the unit cell does indeed double along a symmetry axis when the material transforms into its low-temperature phase. Moreover, as part of our neutron diffraction work on this compound and its deuterated variant, we have studied their structural transformation as a detailed function of temperature near T_{c2} . As described in Section III, our neutron diffraction data confirm that the transformation is first-order (occurring at a slightly lower temperature in the deuterated compound) and that it involves the structural changes revealed by our x-ray measurements.

It is quite conceivable that, at temperatures just above the lower critical point T_{c2} , there would occur a softening of a phonon mode, which anticipates dynamically the static structural change (i.e., the unit-cell doubling) that ultimately takes place at T_{c2} . Although the abruptness of the transition would disallow the mode softening from growing continuously all the way to completion, it was nevertheless thought that at least the starting stages of a phonon instability would be detectable in a sensitive measurement. We have therefore carried out a fairly intense survey of inelastic neutron scattering measurements on LiNH_4SO_4 and LiND_4SO_4 crystals. Our search for soft phonons was ultimately unsuccessful due to an apparent disappearance of phonon scattering near the Brillouin-zone boundary. We nevertheless did obtain some potentially very useful phonon-dispersion information. This work is discussed in Section IV.

II. THE CRYSTAL STRUCTURE OF LiNH_4SO_4 AT ROOM TEMPERATURE

The room-temperature neutron diffraction experiments on LiNH_4SO_4 crystals were performed on the 4-circle diffractometer at the CP-5 reactor of the Argonne National Laboratory, in collaboration with Dr. Selmer W. Peterson of the Argonne staff. The two single-crystal samples used in these experiments were supplied to us by the Philips North American Corporation and were approximately 10 mg and 70 mg in weight. With a monochromatic incident neutron beam, integrated scattering intensity data were collected for over one hundred Bragg reflections, for which background corrections were made on the basis of individual rocking curves. The data taken on the two crystals for the reflections of weak and moderate intensities were in excellent agreement. For the strong reflections, however, reliable intensity information could only be obtained with the smaller crystal for which extinction effects were very small.

Fourier maps made of all our neutron scattering data verified that the crystal structure of LiNH_4SO_4 belongs to the space group $P2_1$ cn with orthorhombic unit cell dimensions, $a = 5.280 \text{ \AA}$, $b = 9.140 \text{ \AA}$, $c = 8.786 \text{ \AA}$, as determined earlier by Dollase from x-ray diffraction measurements.[9] However, our results extended beyond those of the x-ray work in that they were able to locate the positions of the hydrogen atoms. As shown in Fig. 1, the four hydrogens of each NH_4 complex form a slightly distorted tetrahedron with the nitrogen essentially at the geometric center, the N-H distances varying from 0.938 \AA to 0.996 \AA . The H-N-H

angles vary between about 106° and 112° , the ideal tetrahedral angle being 109.5° . The orientation of each NH_4 tetrahedron is such that the hydrogens are placed in close proximity with the nearest oxygens of the surrounding network of SO_4 and LiO_4 tetrahedra. Thus, the existence of strong H-O bonds is clearly manifested in the structure. Specifically, the H-O bonds are oriented such that two of the hydrogen atoms are approximately on the perpendicular-bisecting plane of the line joining neighboring oxygen pairs. Moreover, the thermal vibration ellipsoid for these hydrogen atoms is highly eccentric, with its long axis lying in this plane. The other two hydrogen atoms have only one oxygen neighbor and their thermal vibration ellipsoids are nearly spherical.

The room-temperature crystal structure of LiNH_4SO_4 having thus been determined completely, we proceeded to calculate its spontaneous ferroelectric polarization. For these calculations, we adopted a point-charge model in which the ionic valences were set to be Li^+ , S^{6+} , and O^{2-} , while the NH_4 complexes were considered in several alternative cases. In the first case we assumed H^+ and N^{3-} , and in the second case H^- and N^{5+} . The calculated polarizations were respectively 241.6 and 157.0, in $\mu\text{C}/\text{cm}^2$ directed parallel to the \underline{a} axis (see Fig. 1). We also considered the case of H^0 and N^+ , and obtained the intermediate polarization value of 198.7 $\mu\text{C}/\text{cm}^2$, again parallel to \underline{a} . All three of these

calculated values for the room-temperature spontaneous polarization of LiNH_4SO_4 are in gross disagreement with the experimentally determined values of $\sim 0.3 \mu\text{C}/\text{cm}^2$ [5] and $\sim 0.8 \mu\text{C}/\text{cm}^2$ [7]. However, at least one of these two sets of experimental results may well be suspect since they disagree on the sign of the abrupt polarization change at the 11°C transition. Nevertheless, the enormous discrepancy between the calculated and experimental values for the room-temperature polarization is probably real and most likely arises from our simplistic assumption of a completely ionic model in which no allowance was made for any degree of covalency resulting in some charge distributed in the ligands. Such an allowance, however, would require information about the electronic wavefunctions of LiNH_4SO_4 that is presently unavailable.

III. THE STRUCTURAL NATURE OF THE LOW-TEMPERATURE FERROELECTRIC TRANSITION IN LiNH_4SO_4

X-ray diffraction experiments were undertaken in order to investigate the structural changes that may accompany the abrupt ferroelectric transition in LiNH_4SO_4 at $\sim 11^\circ\text{C}$. This work also was carried out in collaboration with Dr. S. W. Peterson at the Argonne National Laboratory. For these experiments, a small piece taken from the larger crystal used in our neutron diffraction studies was mounted on a cryogenic stage attached to

an x-ray precession camera. For studies below the transition point, the sample temperature was maintained at -1.5°C by means of a steady flow of the cool vapor of liquid nitrogen. At this temperature and at room temperature, precession photographs were taken along the orthorhombic a^* , b^* , and c^* axes in various reciprocal-lattice planes. The results obtained in the $(h0\ell)$ planes are shown superimposed for the two temperatures in Fig. 2. It is clear from this figure that a new set of spot images appeared at -1.5°C , which have one-half the spacing along \vec{c}^* of those observed at room temperature. Specifically, from the numerical values shown for the principal spacings: at -1.5°C the ratio $a^*/c^* \approx 8.1/2.4 \approx 3.37$, whereas at room temperature $a^*/c^* \approx 8.1/4.8 \approx 1.69$. Thus, the unit cell in real space doubles along the c axis (i.e., normal to the polar a axis) when the material is cooled through its ferroelectric transition at $\sim 11^{\circ}\text{C}$. This is precisely the structural change that was suggested earlier on the basis of optical measurements.[8] Furthermore, our results appear to indicate that the orthorhombic symmetry is preserved during this structural change. However, we find that the intensities and absences of the various reflections for the low temperature phase cannot be explained on the basis of any orthorhombic space group. We believe that this difficulty probably arises from the existence of crystallographic twins of presumably lower symmetry. If this is so, very careful work will be needed to suppress the twinning and allow the low temperature structure to be identified.

While carrying out neutron scattering measurements of the phonon dispersion in a LiNH_4SO_4 crystal and a deuterated (LiND_4SO_4) crystal, which will be discussed in the following section, we monitored the intensities of selected Bragg reflections as the temperature was varied through the ferroelectric transition point. In the case of the undeuterated crystal, the (003) reflection was chosen, which is forbidden for the room temperature phase. As shown in Fig. 3, the intensity of this reflection increased abruptly at $\sim 9^\circ\text{C}$ as the temperature was lowered and then decreased with similar abruptness at $\sim 12^\circ\text{C}$ as the temperature was subsequently raised. These data support our x-ray diffraction results for the structural change that accompanies this transition and also show that the change takes place essentially discontinuously with a sizable temperature hysteresis. Thus, the ferroelectric transition in LiNH_4SO_4 is unambiguously first-order, which is consistent with previous reports based on dielectric measurements.[5,7] In the case of the deuterated crystal, both the (200) and (20 1/2) reflections (the latter indexed as (201) for the low temperature phase) were monitored in intensity as the temperature was slowly raised through the critical region. It was observed that while the (200) intensity exhibited virtually no change, the (20 1/2) intensity dropped discontinuously at $\sim 7^\circ\text{C}$, as shown in Fig. 3. These data are also consistent with our x-ray diffraction results, indicating that LiND_4SO_4 and LiNH_4SO_4 undergo the same crystallographic

change at their ferroelectric transition. Furthermore, the transition in LiND_4SO_4 is also first-order but occurs at a slightly lower temperature, which again agrees with the results of previous dielectric measurements.[4]

IV. SEARCH FOR SOFT PHONONS IN LiNH_4SO_4 AND LiND_4SO_4

Inelastic neutron scattering measurements were initially performed on a LiNH_4SO_4 crystal at the CP-5 reactor of the Argonne National Laboratory. Having found that the structural change in LiNH_4SO_4 , when cooled through its ferroelectric transition point, involves a unit-cell doubling along the orthorhombic c axis, our search for a soft phonon mode just above the transition point was concentrated in the reciprocal-space region of the strongest superlattice reflections, such as $(20 \frac{1}{2})$. This region corresponds to that of the Brillouin-zone boundary where the wavevector $\vec{k} = \frac{1}{2} \vec{c}^*$. Thus, energy scans were made, typically up to ~ 8 meV, with the neutron scattering vector \vec{q} parallel to \vec{c}^* and fixed in magnitude at some value between zero and $\frac{1}{2} c^*$ ($=\pi/c$), as measured from the reciprocal lattice point. However, due to an enormous q-dependent background produced by incoherent scattering, inelastic scattering peaks attributable to phonons could only be detected with difficulty.

Since the large incoherent scattering was thought to arise primarily from the hydrogen, the LiNH_4SO_4 crystal sample was replaced by a deuterated (LiND_4SO_4) crystal. The sample replacement was extremely effective in reducing the scattering background, and it immediately became much easier to observe the room-temperature inelastic scattering from a transverse acoustical phonon branch (with polarization parallel to \vec{a}) but only at q values up to $\sim 0.2c^*$. At larger values of q , where this phonon branch extends to higher energies (E), the intensity of the inelastic scattering peak was seen to diminish as $1/E^2$ (as expected theoretically) and thus to rapidly become unmeasurable.

In order to follow this acoustical phonon branch closer to the Brillouin zone boundary and to observe any other phonon modes at higher energies, we needed access to a more intense neutron source. Hence, the LiND_4SO_4 crystal was transferred to the high-flux reactor at the Brookhaven National Laboratory, where the remainder of our inelastic scattering experiments were performed. With an order-of-magnitude increase of incident neutron flux, the transverse acoustical (TA) phonon branch seen at Argonne was followed out to $q \approx 0.3c^*$, again at room temperature. Beyond this value of q , there was an anomalously rapid decrease in the observed intensity of the inelastic scattering peak, making it impossible to learn if this phonon branch ultimately softens (i.e., dips to lower energies) near the Brillouin-zone boundary. Up to $q \approx 0.3c^*$, the phonon dispersion for this TA branch is

linear, as shown in Fig. 4, corresponding to a sound velocity (v_s) of $\sim 2.2 \times 10^5$ cm/sec. Moreover, as indicated in the figure, there was no apparent change in this phonon dispersion curve when the sample temperature was lowered first to 7.5°C and then to -40°C . At the latter temperature, we also observed a longitudinal acoustical phonon branch at higher energies (see Fig. 4), but again the inelastic scattering peak was found to disappear when q reached about half-way toward the zone boundary; in this case, the linear dispersion at low q gives $v_s \approx 4.9 \times 10^5$ cm/sec. There seems to be no reasonable explanation for the disappearance of the scattering from both these phonon branches near the zone boundary; further work is needed to clarify this perplexing result.

REFERENCES

1. J. Kobayashi, Y. Enomoto, and Y. Sato, Phys. Stat. Sol.
(b) 50, 335 (1972).
2. V. Dvorak, Ferroelectrics 7, 1 (1974).
3. W. A. Smith, Philips Laboratories Quarterly Technical Reports,
June-Aug. 1977 and Sept.-Nov. 1977.
4. P. Kumara Acharya and P. S. Narayanan, Indian J. Pure Appl.
Phys. 11, 514 (1973).
5. V. I. Yuzvak, L. I. Zharebtsova, V. B. Shkuryaeva, and I. P.
Aleksandrova, Soviet Phys. Crystallogr. 19, 480 (1975).
6. B. O. Hildmann, Th. Hahn, L. E. Cross, and R. E. Newnham,
Appl. Phys. Lett. 27, 103 (1975).
7. T. Mitsui, T. Oka, Y. Shiroishi, M. Takashige, K. Iio,
and S. Sawada, J. Phys. Soc. Japan 39, 845 (1975).
8. H. Poulet and J. P. Mathieu, Solid State Commun. 21, 421
(1977).
9. W. A. Dollase, Acta Cryst. B 25, 2298 (1969).

FIGURE CAPTIONS

- Fig. 1. The crystal structure of LiNH_4SO_4 at room temperature, shown projected on the a-b plane, where the dotted lines define half the orthorhombic unit cell. The elevations of the atoms are indicated in fractional units of c, except for the Li atoms, which lie approximately $1/2$ c above the S atoms. Only one of the two NH_4 complexes in this half-cell are represented (see Ref. 9).
- Fig. 2. Composite representation of x-ray precession photographs obtained for the $(h0\ell)$ planes of LiNH_4SO_4 at -1.5°C and 20°C . The ℓ values shown are appropriate for the low-temperature phase and are twice those of the room-temperature phase.
- Fig. 3. Neutron Bragg intensities (in arbitrary units) of the (003) reflection from LiNH_4SO_4 and of the $(20\ 1/2)$ reflection from LiND_4SO_4 for increasing or decreasing temperature.
- Fig. 4. Inelastic neutron scattering results for the phonon dispersion (energy in meV vs wavevector parallel to \vec{c}^*) in LiND_4SO_4 , for a transverse acoustical (TA) mode with phonon polarization parallel to \vec{a}^* and for a longitudinal acoustical (LA) mode, at various temperatures.






Figure 1



- image observed only at -1.5°C
- ⊙ image observed at -1.5°C and 20°C

Figure 2




Figure 3




Figure 4

DISTRIBUTION LIST FOR FINAL TECHNICAL REPORTS

(1 Copy Unless Otherwise Specified)

Defense Documentation Center
 Attn: DDC-TCA
 Cameron Station, Bldg. 5
 Alexandria, VA 22314
 (12 copies)

Director
 National Security Agency
 Attn: TDL
 Fort George G. Mead, M.D. 20755

Office of Naval Research
 Code 417
 Arlington, VA 22217

Director
 Naval Research Laboratory
 Attn: Code 2627
 Washington, D. C. 20375

Commander
 Naval Electronics Laboratory Center
 Attn: Library
 San Diego, CA 92152

Commander
 U.S. Naval Surface Weapons Laboratory
 Attn: Dr. F. Bis
 White Oak, Silver Spring, MD 20910

Commandant
 Marine Corps
 HQ, US Marine Corps
 Attn: Code LMC
 Washington, D. C. 20380

Air Force Avionics Laboratory
 Attn: AFAL/TSR, STINFO
 Wright-Patterson Air Force Base
 Ohio 45433

HQDA (DACE-CMS)
 Washington, D.C. 20310

OSASS-RD
 Washington, D. C. 20310

CDR, US Army Material Command
 Attn: AMCMA-EE
 5001 Eisenhower Avenue
 Alexandria, VA 22333

CDR, US Army Materiel Command
 Attn: AMCRD-FW
 5001 Eisenhower Avenue
 Alexandria, VA 22333

Commander
 US Army Missile Command
 Attn: AMSMI-RR
 Dr. J. P. Hallows
 Redstone Arsenal, AL 35809

Commander
 US Army Armament Command
 Attn: AMSAR-RDP (Library)
 Rock Island, IL 61201

Commander
 Picatinny Arsenal
 Attn: SARPA-TS-S #59
 Dover, NJ 07801

Commander
 Frankford Arsenal
 Attn: Dr. William McNeill PDS
 Philadelphia, PA 19137

Commander
 USASA Test & Evaluation Center
 Fort Huachuca, AZ 85613

US Army Research Office-Durham
 Attn: CRDARD-IP
 Box CM, Duke Station
 Durham, NC 27706

US Army Research Office-Durham
 Attn: Dr. Robert J. Lontz
 Box CM, Duke Station
 Durham, NC 27706

Aerojet
Electro System Company
Attn: P. C. Wang
1100 W. Hollyvale Street
Azusa, CA 91702

Arthur D. Little Incorporated
Acron Park
Attn: Jacques M. Steininger
Cambridge, MA 02140

Barnes Engineering
Attn: Dr. W. Rolls
30 Commerce Road
Stamford, CT 06902

Battelle Memorial Institute Library
505 King Avenue
Columbus, OH 43201

Cincinnati Electronics Corporation
Attn: Dr. Vernon Lambert
2630 Glendale-Milford Road
Cincinnati, OH 45241

Electro-Optical Systems, Inc.
Solid State Department
Attn: Dr. P. J. A. Zoutendyk
300 N. Halstead Street
Pasadena, CA 91107

Fairchild Company
Charge Coupled Device Department
Attn: Dr. R. H. Dyck
Palo Alto, CA 94304

Ford Motor Company
Scientific Research Staff
Attn: Dr. H. Holloway
P. O. Box 2053
Dearborn, MI 48121

General Dynamics
Attn: Dr. Ester Krikorian
Mission Blvd.
Pomona, CA 91766

Honeywell Inc.
Aerospace & Defense Group
Infrared Detector Group
Attn: George D. Anderson
1611 North Kent Street
Arlington, VA 22209

Hughes Research Laboratories
Attn: Lloyd DeVaux
3011 Malibu Canyon Road
Malibu, CA 90265

RCA Laboratories
Attn: R. L. Foley
Princeton, N. J. 08540

Lincoln Laboratory
Massachusetts Institute of Tech.
Attn: Dr. I. Melngailis
Lexington, MA 02173

Perkin-Elmer Corporation
Associate Director of Research
Attn: Dr. David A. Huchital
Norwalk, CT 06852

Aeroneutronics Ford Corporation
Ford Road
Attn: Mr. Sutton
Newport Beach, CA 92663

Rockwell International Corporation
Autonetics Division
Attn: Dr. G. Hover or
Dr. R. C. Geiss
P. O. Box 4192
3370 Miraloma Avenue
Anaheim, CA 92803

Rockwell International Corporation
Science Center
Attn: Library (H. M. Coogan)
1049 Camino Dos Rios
Thousand Oaks, CA 91360

Globe-Union Inc.
5757 N. Green Bay Ave.
Milwaukee, Wisc 53201

Commander
 HQ MASSTER
 Technical Information Center
 Attn: Mrs. Ruth Reynolds
 Fort Hood, TX 76544

USA Security Agency
 Attn: IARD
 Arlington Hall Station
 Arlington, VA 22212

Commander
 U. S. Army Missile Command
 Attn: AMSMI-RE (Mr. Pittman)
 Redstone Arsenal, AL 35809

Commander
 US Army Systems Analysis Agency
 Attn: (Mr. A. Reid), AMXSY-T
 Aberdeen Proving Ground, MD 21005

NASA Scientific & Tech Info Facility
 Attn: Acquisitions Branch (S-AK/DL)
 P. O. Box 33
 College Park, MD 20740

Chief
 Ofc of Missile Electronic Warfare
 Electronic Warfare Lab, ECOM
 White Sands Missile Range, NM 88002

Director
 Defense Advanced Research
 Projects Agency
 1400 Wilson Blvd.
 Attn: Dr. R. Reynolds
 Arlington, VA 22209
 (3 copies)

Naval Postgraduate School
 Attn: Dr. F. Tao (52-TV)
 Monterey, CA 93940

USNADC
 Attn: Dr. M. Hess
 Code 202149
 Warminster, PA 18974

Institute for Defense Analyses
 Attn: Dr. A. D. Schnitzler
 400 Army-Navy Drive
 Arlington, VA 22202

Infrared Information and
 Analysis Center
 P. O. Box 618
 Ann Arbor, MI 48107

Advisory Group on Electron Devices
 201 Varick Street, 9th Floor
 New York, NY 10014

Commander
 US Army Electronics Command
 Attn: DRSEL-MS-TI
 Fort Monmouth, N. J. 07703

Commander
 US Army Electronics Command
 Attn: DRSEL-PL-ST
 Fort Monmouth, NJ 07703

Aeronutronics Ford Corporation
Aeronutronic Division
Attn: Mr. J. Roschen
Ford Road
Newport Beach, CA 92663

Honeywell Corporate Research Center
Attn: Dr. D. Long C-1300
10701 Lyndale Ave. South
Bloomington, NM 55420

Honeywell Incorporated
Attn: Dr. N. Reine
2 Forbes Road
Lexington, MA 02173

Texas Instruments Incorporated
Central Research Laboratories
Attn: Dr. Chapman
P. O. Box 5936
13500 North Central Expressway
Dallas, TX 75222

S. Campana/Code AEYE
Naval Air Development Center
Warminster, PA 18974

K. J. Ando
Jet Propulsion Laboratory
California Institute of Technology
4800 Oak Grove Drive
Mail Stop 168-222
Pasadena, CA 91103

D. F. Barbe
Code 5214
Naval Research Laboratory
Washington, D. C. 20390

R. A. Belt
USAD Avionics Laboratory
AFAL/TEO-3 Bldg. 620
Wright-Patterson Air Force Base, OH 45433

L. W. Sumney
Code 0335
Naval Electronics System Command
Washington, D. C.

General Electric Company
Attn: Mr. David Schmidt
7777 Leesburg Pike
Falls Church, VA 22043

Hughes Aircraft Company
Aerospace Group
Attn: Dr. K. Nummedall
Bldg 5, Mail Station B 169
Culver City, CA 90230

Raytheon Company
Special Micro-wave Devices
Operation
Infrared and Optical Research
Laboratory
Attn: Dr. P. Debye
130 Second Avenue
Waltham, MA 02154

Phillips Laboratories
345 Scarborough Road
Briarcliff Manor, NY 10510

Westinghouse Electric Corporation
Research & Development Center
Attn: Dr. H. C. Nathanson
Pittsburgh, PA 15235

Rockwell International
Autonetics Division
Attn: R. Aquilera
Anaheim, CA 92803

Stanley Sobieski
Code 672
Goddard Space Flight Center
Greenbelt, MD 20771

Technology Associates
Attn: Dr. O. Trapp
51 Hillbrook Drive

Portola Valley, CA 94025

Air Force Materials Laboratory
Attn: Robert Spry
LPO Wright-Patterson
Air Force Base, OH 45433